

Wednesday, 13 September 2017

## Nursing



The image shows a poster titled "FOLIC ACID and VITAMIN D GUIDELINES FOR HEALTH PROFESSIONALS". It is produced by the HSC Public Health Agency and the HSC Health and Social Care Board. The poster is divided into several sections:

- FOLIC ACID AND NEURAL TUBE DEFECTS (NTD)**
  - Baseline for ALL women who could become pregnant:**
    - Advise women that they can purchase Folic Acid (400 mcg daily) over the counter. OR
    - Prescribe 400mcg daily Folic Acid
  - Duration:** from 12 weeks before conception to 12 weeks pregnant
- A woman is AT HIGHER RISK of having a baby with a Neural Tube Defect (NTD) affected pregnancy if one or more of the following apply:**
  - There is a family history of NTDs (woman or father's family)
  - She has diabetes
  - She is taking anti-epileptic drugs\* (for any indication)
  - She has a BMI > 30
  - She has coeliac disease
  - She has thalassaemia
- For women AT HIGHER RISK of having a baby with an NTD:**
  - Prescribe 8mg daily Folic Acid (prescription-only dosage)
  - Duration: from 12 weeks before conception to 12 weeks pregnant
- VITAMIN D – RDA 10mcg a day**
  - In summer, most people get enough vitamin D from sunlight. Between October and March a 10mcg supplement of vitamin D per day for pregnant/breastfeeding women should be encouraged. [People who cover their skin for cultural reasons, those confined indoors or with dark skin should take a daily supplement of vitamin D throughout the year.]

Folic acid and Vitamin D guidelines for health professionals.

## Details

Format

PDF, poster

Target group

Health Professionals

## Downloads

Attachment	Size
<a href="#">FOLIC ACID and VITAMIN D Guidelines 2017.pdf</a>	170.75 KB

## Tags

- [folic acid](#)
- [vitamin d](#)
- [pregnancy](#)
- [nutrition](#)
- [supplements](#)

[Print](#)