Tuesday, 30 April 2019

Health Protection

HSC Public Health Agency

Clostridium difficile

Clostridium difficile, also known as C. difficile or "C. diff", is bacteria that can infect the bowel and cause diarrhoea

The infection most commonly affects people who have recently been treated with antibiotics. It can spread easily to others.

C. difficile infections are unpleasant and can sometimes cause serious bowel problems, but they can usually be treated with another course of antibiotics.

Symptoms of a C. difficile infection

Symptoms of a C. dtfficile infection usually develop when you're taking antibiotics, or when you have finished taking them within the last few weeks.

The most common symptoms are:

- · diarrhoea several times a day
- fever · loss of appetite
- feeling sick
 tummy pain

In some cases, you may have signs of dehydration.

How you get C. difficile

C. difficile bacteria are found in the digestive system of about 1 in every 30 healthy

The bacteria often live hamilessly because other bacteria normally found in the bowel keep it under control

But some antibiotics can interfere with the balance of bacteria in the bowel, which can cause the C difficile bacteria to multiply and produce toxins that make the person il

When this happens, C. difficile can spread easily to other people because the bacteria are passed out of the body in the person's diarrhoea.

Once out of the body, the bacteria turn into resistant cells called spores.

These can survive for long periods on hands, surfaces (such as toilets), objects and clothing unless they're thoroughly cleaned, and can infect someone else if they get into their mouth.

This leaflet gives info on Clostridium difficile, also known as C. difficile or "C. diff", a bacteria that can infect the bowel and cause diarrhoea.

Details

Format A4 Pdf Target group Health Professionals and general public

Downloads

Attachment Size Clostridium difficile factsheet.pdf 269.98 KB

<u>Print</u>