


Wednesday, 02 November 2022

## Health Protection



**A guide to COVID-19 vaccine doses, boosters and interval between doses**

**COVID-19 vaccine**

**Primary course**

The **primary course** describes the vaccines given to individuals who have never received the vaccine before. The purpose of the primary course is to provide **new** protection (immunity) for the individual. For the COVID-19 vaccine:

- a complete primary course is two primary doses either 8 or 12 weeks apart – this interval between doses depends on age and whether someone is immunosuppressed or has a weakened immune system;
- a small number of people with certain medical conditions also require a third primary dose to complete the 'primary course'. People who fall into this category will be notified of this by their hospital clinician or GP.

**Booster dose**

For some vaccines, protection (immunity) declines over time. A **booster dose** is the name for vaccines given to individuals who have completed the primary course and need to have their existing protection (immunity) increased or 'boosted'.

The number and frequency of booster doses an individual requires depends on their age and underlying health.

For the COVID-19 vaccine:

- individuals aged 16 years and over who completed their primary course (whether a 2 dose or 3 dose course) should have a booster dose at least 3 months after their last dose;
- those individuals who are aged 75 and over, resident in a care home or aged 12 and over and are immunosuppressed and received their last dose at least 3 months ago should have a spring booster dose from April 2022;
- planning is taking place for an autumn booster programme for eligible individuals.

Vaccine doses have been named and summarised below to help improve understanding of the different doses:

Vaccine dose	Description
1st primary dose	2 dose primary course.
2nd primary dose	
3rd primary dose	People with certain medical conditions will not respond as well to 1st and 2nd primary doses, and they should have a 3rd primary dose to complete the 'primary course' of vaccination.
Booster dose	Individuals aged 16 years and over who completed their primary course (whether a 2 dose or 3 dose course) should have a booster dose at least 3 months after their last dose.
Spring 2022 booster programme	This booster dose is only for specific high risk groups; those aged 75 years and over, care home residents and immunosuppressed people aged 12 years and over.
Autumn 2022 booster programme	This booster programme is being planned for autumn and the eligible groups are being decided.

HSC Public Health Agency  
Public Health Agency  
12-22 Lincolns Street, Belfast BT3 9SS.  
Tel: 0950 535 0114 Social media.  
www.publichealth.hscni.net

Find us on: Facebook, Twitter, Instagram, YouTube

COVID-19

This resource explains the COVID-19 vaccine doses, booster doses and interval between doses for each eligible group.

## Details

Format

A4 1 side only

Target group

Health professionals and vaccinators

## Downloads

**Attachment**

**Size**

[A guide to COVID-19 vaccine doses Final.pdf](#) 461.35 KB

## Tags

- [covid-19 vaccine](#)

[Print](#)