

Children's Bladder and Bowel Care

Level 1 Resource Pack



INDEX

Introduction.....	page 3
Assessing parental intolerance.....	page 3
Night time wetting initial advice.....	page 4
Fluid advice	page 5
Strategies to help children increase their fluid intake	page 6
Dietary advice.....	page 7
Symptoms of constipation.....	page 7
Toileting advice.....	page 7
Skills for toilet training.....	page 9
Children who give no indication of needing to use the toilet.....	page 10
Toileting assessment.....	page 11
Toilet skills assessment tool	page 15
Toileting chart.....	page 17
Frequency volume chart	page 19
Children’s bladder and bowel initial assessment tool.....	page 21
References and resources.....	page 24
Resources for professionals.....	page 24
Resources for families	page 24

Introduction

Bladder and bowel issues have a huge impact. They cause shame, embarrassment and frustration, as well as social isolation, feelings of difference, reduced self-esteem and social opportunities and lower quality of life for the whole family. Furthermore, for some children and young people, they affect sleep, which may impact on behaviours and learning in the day and for others they may result in reduced school attendance.

This resource pack includes the information to help healthcare professionals working within primary care settings to support children and young people who present with bladder and bowel issues, including delayed toilet training, constipation, soiling, lower urinary tract symptoms including frequency, urgency, day or night time wetting.

Further advice and support should be available from the local specialist children's bladder and bowel service.

Assessing parental tolerance

Parents or carers who are intolerant of their child will focus on the impact on themselves rather than on the child and may use punishment inappropriately. This may result in concerns for the child's welfare.

Questions to ask include:

What concerns you about the wetting/soiling? Supportive parents or carers will express concern for the emotional state and wellbeing of their child, impact on their child's social activities and on their self-esteem. Parents or carers who may be intolerant are more likely to focus on the impact of extra washing and drying, the smell and the cost of replacing bedding or clothing.

What are the reasons for the wetting/soiling? Supportive parents or carers may link incontinence to causes outside their child's control, such as deep sleep or family history. Intolerant parents may consider their child to be lazy, doing it on purpose, or doing it to get back at or punish the parents in some way.

What has your child tried to do to stop the wetting/soiling? Supportive parents or carers may talk about attempts made by their child, such as helping to get changed, following instructions from the parents, e.g. drinking more or less, stopping fizzy drinks. Intolerant parents or carers are more likely to consider their child can be dry or clean when they want to be, not to be bothered, or having not tried anything.

How does the wetting/soiling make the parent or carer feel? Parents or carers who are supportive of their child may talk about being empathetic with them and how it is unpleasant for their child. Intolerant parents or carers may express hostility, anger, annoyance or frustration with their child.

How do you cope with the wetting/soiling? Parents or carers who are supportive try to find solutions and cope with practicalities. Those who are intolerant may be punishing their

child, humiliating them, showing disappointment, making threats, reprimanding or withdrawing privileges.

(Adapted from Nocturnal Enuresis Resource Pack, Charts Questionnaires and Information to Assist Professionals, R Butler, fifth edition, 2006. Pub: ERIC)

It is not unusual for parents and carers to consider the impact on themselves as well as their child. However, the clinician should be observant for any signs that may suggest that there are safeguarding concerns. If these are present, appropriate and timely advice should be sought from the Children's Bladder and Bowel Service, and/or from the safeguarding children department with action taken and documented in accordance with the local policies and procedures on safeguarding children.

Promoting Continence Pathway: NIGHT TIME WETTING

INITIAL ADVICE

If bedwetting has only started in the last few days or weeks consider whether it might be caused by systemic illness.

If the child also has daytime symptoms (e.g. urgency, frequency, daytime wetting), refer to the daytime wetting pathway.

- Explain that the bedwetting is not deliberate, nor is it the child's fault and that the child should not be told off or punished.
- If possible, explain the causes of bedwetting.
- Encourage daytime water-based drinks (see fluid advice). Avoid caffeinated, fizzy and energy drinks.
- Do not limit fluid intake during the day, unless excessive (see fluid advice).
- Encourage regular daytime toileting (about two hourly).
- Encourage the child to try and pass urine before settling for sleep each night.
- Advise that the child should avoid all food and drink in the last hour before sleep.
- Avoid high salt and high protein foods late in the day (these increase urine production)
- If the child is using products (e.g. pyjama pants, nappies) and family circumstances allow, consider a trial of at least two consecutive nights without.
- Do not lift/wake the child when parents/carers go to bed. The only times when lifting may be acceptable is in the short term when it is particularly important that the bed stays dry e.g. when on holiday.
- If the child wakes themselves during the night, ask parents/carers to encourage them to use the toilet before settling back to sleep.
- Discuss ways of reducing the impact of the wetting, such as bed protection, washable or disposable products. E.g. a waterproof sheet on the mattress, or absorbent pants.
- Consider access to the toilet at night. If this is difficult, try to find ways to make it easier e.g. torch by the bed or potty in the room.
- Consider whether the child can get out of bed or has anxieties or fears that may result in difficulties getting up e.g. fear of the dark.

- Advise parents/carers to only use rewards for things that are in their child's control. Remember that a child cannot control what happens when they are asleep. Therefore, encouragement and positive comments should be made for dry nights, but rewards (if used) should focus on things that are in the child's control, such as drinking recommended levels and toileting during the day, for toileting before sleep, helping to strip their own bed etc. Do not give rewards for dry nights but do notice and praise any efforts made by the child.
- Monitor progress by keeping a diary of wet and dry nights, of waking after wetting, of waking to use the toilet.

If there is no improvement with initial advice within two to four weeks consider first line treatment options such as desmopressin or an alarm or refer to the children's bladder and bowel service. There is evidence that persisting with lifestyle adjustments alone, is not helpful in overcoming bedwetting.

FLUID ADVICE

Adequate fluid intake is an important part of treatment for bladder and problems affecting children and young people, including daytime wetting, night time wetting and constipation. It is also important when children are learning the skills for toilet training.

- Caffeinated drinks, including tea, coffee, hot chocolate, energy drinks and cola, should be avoided as they may have a diuretic effect and can contribute to bladder overactivity.
- Fizzy drinks should be avoided as they can contribute to bladder overactivity.
- Children and young people need extra water-based fluids if they are doing lots of exercise (including sports, playing out and school playtimes), or if the weather or their environment is hot.
- Milk is healthy, but is used by the body as a food. It should not be encouraged instead of, or as part of total water-based drinks.
- Do not restrict fluid intake. If fluid intake is excessive, consider whether this may be due to a sensory issue, behavioural issues or if the child / young person may have diabetes insipidus.
- Children and young people should be encouraged to take full water bottles (500 – 750mls) to school and drink the contents.
- Water is the healthiest drink and should be encouraged. However, many children refuse to drink it. If children / young people do not like to drink water, arrangements should be made for them to take a non-see-through bottle to school with diluted fruit squash (preferably sugar-free).
- Schools should be asked to allow the child open access to their drinks bottles and to the toilet, particularly if they are being encouraged to increase their drinking, or have day time bladder or bowel issues.

Age	Sex	Total drinks per day
7-12 months		600 – 900ml
1-3 years	Female	900 – 1000ml
	Male	900 – 1000ml
4–8 years	Female	1200–1400 ml
	Male	1200–1400 ml
9–13 years	Female	1200–2100 ml
	Male	1400–2300 ml
14–18 years	Female	1400–2500 ml
	Male	2100–3200 ml

Suggested intake of water based drinks per 24 hours according to age and sex:

(Adapted from CG 111 Nocturnal Enuresis NICE 2010 and American dietary requirements, cited in CG 99 Constipation in Children and Young People, NICE 2010)

NB higher intakes of water are required when children are physically active, or the weather or environment is hot.

Overweight children may also require more water.

Strategies to help children increase their fluid intake:

- Positive reinforcement for drinking well, including use of appropriate charts and rewards.
- Start with expecting the child to drink only slightly more than they currently are and gradually increase expectations.
- Measure out what the child should be having each day into a clean jug or plastic bottle. Making all their drinks from that can help them visualise how well they are doing. If they have a drink from a carton or bottle, the equivalent quantity of water from the jug or bottle can be poured away. The child should be encouraged to finish their jug or bottle by the end of teatime each day.
- Some children manage well if given a full glass and are told to drink half, others do better if given half a glass and are told to finish it.
- Build drink times into the family’s daily routine.
- Make drink times fun: Suggest that the parent or carer sit together with their child and read a book or play game and do not read any more/throw the dice until the child has had a few more sips. If the child refuses to drink more the parent can put the book or game away.
- Use straws or a different glass or cup, with a design that is appealing to the child.
- Add ice or give the child the drink at the temperature they prefer.
- If children really dislike or refuse to drink water, offer fruit squashes or diluted fruit juices. Many of these are high in sugar content – sugar free alternatives are best, for children who will not drink water.
- Ice lollies and jellies are high in fluid content, but tend to be high in sugar, so should be used with caution.
- Families should be advised to avoid battles over drinks.
- The child should be having half their daily intake by dinner time (midday meal) to avoid them having large quantities late in the day, as this may cause or exacerbate night time wetting.

DIETARY ADVICE

- Dietary adjustment alone is not an acceptable treatment for chronic constipation in children and young people, but plays a part in treatment and is part of health promotion.
- Children/young people should be encouraged to eat five or more portions of fruit and vegetables per day.
- Children/ young people should not be encouraged to eat large amounts of high fibre foods (such as wheat biscuits or all bran) as this can exacerbate constipation if fluid intake is inadequate .
- Children / young people should not be eating unprocessed bran.
- Wholegrain cereals, brown bread and rice can be helpful and are part of a healthy diet.
- Children over the age of one year should not be having more than a pint of milk or its equivalent (yoghurts, fromage frais, cheese, custards, rice puddings etc) per day. This can exacerbate constipation, reduce appetite and prevent children from having a balanced, varied diet.

N.B. Please always follow any advice from the dietitian and ensure that the child does not have any foods to which they may have intolerances or allergies.

SYMPTOMS OF CONSTIPATION

Constipation in childhood is a common problem. For many it lasts only a few days, but it can become chronic in up to a third of children and is a common reason for referral to secondary care. Chronic constipation is usually idiopathic (it happens spontaneously and/or the cause is not known). Symptoms vary between children. It is possible for children to be having a bowel motion most days, but to be constipated if they are only partially emptying the rectum. Other children may only pass loose stools, which means it is not always easy to diagnose constipation in children.

Two or more of the following list may be indicative of constipation in children:

- Infrequent bowel motions (less than three times / week in children over 3 years)
- Soiling or overflow: offensive smelling stools that are usually passed without the child being aware of them
- Small, hard or very large stools (rabbit droppings or stools that block the toilet)
- Poor appetite, often improves after a large bowel motion
- Abdominal pain that improves after a bowel motion
- Withholding or appearing to strain
- Anal pain
- Painful bowel motions
- Bleeding associated with hard stools
- Previous constipation
- Current or previous anal fissure

The following may also be symptoms of constipation:

- Unpleasant smelling wind or bowels motions

- Excessive flatulence
- Varying texture to bowel motions
- Bowel motions in sleep in children over a year in age
- Abdominal distension
- Lethargy
- Unhappiness, anger or irritability that improves after a large bowel motion

NB. If the child is presenting as acutely unwell, has faltering growth or gross abdominal distension they should be reviewed by the GP or a paediatrician.

TOILETING ADVICE

- Encourage the child / young person to use the toilet regularly during the day. About two hourly is the correct interval for most. However, if the child / young person is wet more often than this, the interval should be shorter to try and ensure that they remain dry.
- Suggest that the child / young person uses the toilet after they have had a drink. When the child is toileting two hourly this can help with fluid intake as well.
- Ensure the toilet is easy to access, clean and well stocked with toilet paper etc. This is particularly important at school. Secondary school children may benefit from a toilet pass. Primary school children may need the teacher to know about the bladder or bowel issue. Having a signal for the child to indicate to the teacher when they need the toilet and them being allowed to leave the classroom promptly may be helpful, rather than them having to wait to ask to go.
- Ensure that smaller children have an insert seat and stool, so they can sit comfortably, with their bottom well, supported, their feet flat on a firm surface and their knees higher than their hips.
- Ensure that children with mobility difficulties or sensory issues have been referred to an occupational therapist for assessment of their toileting needs.
- If the child / young person feel they need to pass urine urgently or suddenly, they may be encouraged to count to five and if the feeling goes away to wait until the next planned toilet visit. If the feeling remains or they are likely to wet if they do not toilet quickly, then they should go straight to the toilet. Open access should be arranged for the toilet at school.
- There is no evidence of benefit from trying to put off passing urine for longer than a few seconds if a child has urgency or daytime wetting and this should not be encouraged.
- Children should be encouraged to remain at the toilet long enough to complete voiding.
- Children should be encouraged to sit on the toilet long enough to complete a bowel action. They should be able to sit privately. For children with constipation and soiling, there is often benefit from allowing them access to the disabled toilet in school as this is often more private than the main toilets.
- If the child is wetting /soiling at school it would be helpful to them to have spare clothes, wipes and plastic bags for the damp clothes, in their bag to allow changing as needed. Parents/carers should provide these from home and a system arranged so that they know when replacements are required.

- Children should be supported to learn to change independently as soon as they have the developmental skills, which is usually from about four years old. If they are wetting/soiling in school, they may need support with learning to change themselves initially, or until their dexterity is sufficiently good for them to manage alone.
- It is not acceptable for schools and nurseries to request parents or carers attend to support their children with personal hygiene.

[Guidance is available from the Department of Education](#) schools to help them manage children with medical needs. There is also Guidance [on Managing Bladder and Bowel Issues in Nurseries, Schools and Colleges](#) on the Bladder & Bowel UK website.

SKILLS FOR TOILET TRAINING

Learning the skills required for toilet training and becoming independent with this should be among the earliest self-care abilities developed by children and is one of the most important. In traditional societies, most children have attained continence by their second birthdays. In the Western World, the age of introduction of skills for toilet training has increased in recent decades. However, children should start to develop the skills for toilet training in the first year of life.

Supporting children to learn the skills for toilet training should not be delayed due to disability. Failure to offer assistance and advice to families to enable children with disabilities to develop the skills required for toilet training may be considered discriminatory. When children remain reliant on continence containment products, rather than toilet training, they may fail to achieve their potential, with associated increased dependence compared to their peers, reduced self-esteem and self-confidence, increased likelihood of abuse and there is more stress for them and their family.

The following may be helpful:

- Ensure the child has a varied diet and adequate fluid intake if possible (see relevant sections of this document).
- Encourage the child to sit on the toilet or potty regularly. The potty may be better for smaller children as they may feel more secure and their feet will be flat and well supported on the floor, their knees will be bent and higher than their hips. However, if the toilet is used, smaller children must have an insert seat and stool on which they can rest their feet when sitting, to ensure they are in the correct position to pass urine or open their bowels.
- If the child has any physical or sensory issues, they should have an early referral to the occupational therapist for assessment of their toileting needs. Information on [the impact of sensory issues on toilet training](#) is available on the Bladder & Bowel UK website.
- Start by sitting the child on a potty, or adapted toilet once a day for short periods of time and gradually increase frequency and time of sitting. Do not sit the child for more than 3-4 minutes.
- Encourage regular drinks (about 2 hourly) and then potty/toilet times after drinks. About 10-15 minutes later is often best if parents/carers can manage this, otherwise straight away.

- Tip solid poos down the toilet and then flush them away, with the child present.
- Change all nappies in the bathroom.
- If the child is mobile, ensure they are standing to have their nappy changed. Encourage them to be as involved as possible and tip any solid matter down the toilet, involving the child in flushing the toilet.
- Have an open-door policy for toileting so the child sees parents/carers and siblings using the toilet.
- Ensure all carers use the same words to describe wee and poo. Avoid using the word 'dirty' for poo as this has other meanings.
- Discuss the difference between wet and dry.
- Consider using stories, videos etc.
- Children who have communication, processing or learning difficulties are often helped by picture cue cards.
- Encourage the child to learn to help dress and undress themselves.
- Use clothes that are easy for the child to manage.
- Encourage the child to say (or sign) when they are wet or have opened their bowels.
- Modern nappies are very efficient at drawing moisture away from the skin, so reduce the child's awareness of passing urine, or opening their bowels. Consider using cotton underwear or kitchen towel inside the nappy to raise awareness of when they are wet or dry.
- Ensure the parent/carer has a plan for dealing with wetting or soiling when away from home and has good routines established.
- Ask the family to keep an hourly record (see page 16), during their child's waking hours for three or four days of when their child drinks, passes urine or opens their bowels. Use the chart to inform timing of potty or toilet visits, but advise the family not to take their child to the toilet more often than once every 60 minutes.
- When the family are catching about half of the voids in the potty or toilet remove the nappies during the day and continue with timed toilet visits. If the child is wet between visits, the family may reduce the interval between visits, but should not take their child more than once every 60 minutes. This interval can be gradually increased as their child becomes dry between visits, until they are able to go about two hours.
- Start using underwear or training pants during the day.
- Praise and reward success, change in the bathroom when needed with minimum fuss and feedback.
- Consistency is important and once progress is being made, the parent/carer should be encouraged not to return to nappies during the day. Many children will have lots of little voids in the first few days after removal of nappies. This usually improves as the children learn to recognise the bladder signals and to empty their bladder fully on the toilet.

Children who give no indication of needing to use the toilet:

- Consider using cotton underwear or kitchen towel inside the nappy to raise awareness of when they are wet or dry.

- It is only when children start to experience the feeling of being wet that some will become aware of when they need to go to the toilet.
- Consider removing nappies in the day when about half the voids are being caught on the potty or toilet. Parents/carers will need to have a strategy for managing on trips out.
- Regular drinks, (about two hourly) followed by regular trips to the toilet/potty can help the child by ensuring they are voiding more often during the day.
- Consider repeating the record, using a toileting chart (see page 16) of when the child is drinking and when they are passing urine for at least three days, as this can help parents/carers to see their child's natural pattern and help them to get their child to the potty/toilet at the right time.
- Doing all changing in the toilet or bathroom, flushing solid stools down the toilet and sitting the child on the toilet when changing them can all be helpful.
- Using positive reinforcement (praise, reward charts with time-based rewards) for targeted behaviours.
- Many children with learning disabilities or processing differences will take time to learn to indicate when they need the toilet, some may not be able to indicate but will be able to become continent if taken regularly. Giving children options for how they alert their parents/carers is helpful e.g. signing, picture cue cards, consistent word etc.
- Remind parents/carers that their child needs to learn to recognise the bladder/bowel signals that it needs to empty and then how to let someone know, if they are unable to access the potty/toilet independently.
- For children with disabilities, a wetting alarm (the same as a body-worn enuresis alarm) may be helpful. These can usually be borrowed from the children's bladder and bowel service. Provide the family with [instructions](#) on how to use them.

TOILETING ASSESSMENT

Information about skill development for toilet training should be provided to families in the first year of life with assessment for toileting commencing early in the child's second year when there is a physical disability of learning difficulty, or as soon as it is identified that there is any disability or condition that may delay toilet training. Assessment should be dynamic process, with a programme put in place to address any issues. The child should be reassessed every one to three months (depending on individual needs), with the family given an individualised programme to follow in the meantime. The amount of support required will depend on the child's needs and the family dynamics, with some families needing frequent review and others minimal intervention.

The first stage of a toileting assessment involves asking the family to keep a full toileting diary for at least three days using the toileting chart (page 16). This is important as part of promoting bladder and bowel health, even for children who are unlikely to ever be able to toilet train due to the extent of their disability. Failure to fully assess a child's bladder and bowel health may result in problems being missed, with serious long-term consequences. Any problems detected on assessment, such as constipation, constant dribbling of urine, inability to sit, behaviour problems etc, must be addressed.

- The toileting chart should be printed so that the parents/carers have a copy of the instructions that are on the reverse.
- Parents/carers should be advised to keep records for all the child's waking hours for at least three full days. This must be done on days when the child is with the parent/carer all day (i.e. not on school days) and these days do not need to be consecutive. They should also keep records of the child's bowel motions for at least seven consecutive days.
- As modern disposable nappies are so absorbent, it is sometimes difficult to tell the child has voided if they have only passed small amounts of urine. Therefore it is recommended that the child wear cotton pants inside the nappy, or that the parent/carer fold a piece of kitchen towel inside the nappy, as it is obvious when these are wet. The pants or piece of kitchen towel should be changed if they are wet when the nappy is checked, but the nappy does not need to be changed more often than usual.
- The toileting chart should be reviewed when completed to see if:
 - the child is having the recommended intake of drinks,
 - to ensure they are not having excessive milk,
 - to see whether they appear to be having normal bowel actions and
 - to see if they are able to stay dry for more than an hour at a time: if the child is wet every hour the family should be asked to check their child every ten to fifteen minutes for one to two hours. If they are still wet every time they are checked they should be referred to paediatrics to exclude an underlying congenital anomaly. If they are dry on some occasions, then the bladder and bowel service should be asked for advice.
- As promotion of bladder and bowel health is the priority for all children, families should be offered advice as appropriate from the information received from the toileting chart.
- Where a dietician is involved, they should be consulted prior to advice being given to the family about diet, fluid or milk intake.

Once the toileting chart is completed and returned the assessment tool for toilet skills assessment chart (see page 14) must be completed. This should be done with the child and parent/carer, so that the child can be observed in their normal environment, the parent/carer is involved and advice is given in an appropriate and timely way. Carrying out the assessment will allow skills that need additional support to be identified, alongside any underlying pathology. The assessment tool can then be used to inform an individualised toilet skill development programme.

- Sections a) and b) of the toilet skills assessment must be completed using the toileting charts and information observed by the assessor. Normal formed bowel movements (section (b) 2 and 3) refer to a child passing type 3 -5 stools three times a day to once every three days. Any bowel or bladder problem should be addressed using the relevant pathway or discussed with the bladder and bowel service.
- Products are not normally provided for children with enuresis (night time wetting – see section (c)), as this is considered a treatable condition. If the child is dry during the day, the enuresis pathway should be followed.

- Overnight bowel motions in a child who is more than one year old (section d) is normally an indication of constipation. The constipation pathway should be followed.
- Low scores for the section titled Independence (sections (e), (f), and (g)) do not mean that a child cannot toilet train. Efforts should be made to address the problems:
 - If a child is not sitting, then this should be gradually introduced using incentives and encouragement.
 - If a child is not giving any indication of needing to go to the toilet, then sign language, or picture communication may need to be introduced. Individual advice may be sought from the bladder and bowel service.
 - Inability to handle clothes is not a reason for a child to be prevented from toilet training. Assistance should be given to help the child to learn to handle their clothes, where possible. Advice should be provided to the family about using clothes that are easier to adjust, or about appropriate adaptations. The occupational therapist may be able to make suggestions or offer help.
- If it is found that a child never passes urine or opens their bowels on the toilet or potty (sections (h) and (i)) then appropriately timed toileting should be tried. The toileting chart can be used to see if there is any pattern to wetting/soiling, including if these are related to drinks or meals. This information can be used to inform toilet visits. A daytime wetting alarm may increase the child's awareness of when they are voiding.
- High scores for section behaviour problem (section (j)) does not mean that a child cannot learn the skills to toilet train. Efforts should be made to address the problems. Learning disability services may be able to offer some suggestions.
- If a child is likely to require toileting aids or adaptations (section (l)), that should be addressed early and may require referral to the occupational therapist.
- If a child is not responding to basic commands (section (m)), then changing routines or introducing picture cue cards or social stories may be helpful.
- Diet (section (n)) and fluids (section (o)) should be assessed and any changes required discussed with the family, paying heed to individual children's needs or advice given by a dietician if involved.

The toilet skills assessment should be reviewed and actions should be taken as indicated by the prompts. If these actions are felt to be inappropriate this should be documented with the reasons in the child's notes. It is not acceptable to ignore highlighted problems. These must be treated where possible and the child then reassessed.

A individualised toilet training programme should be put in place to support the child develop the required skills. This should include:

- Teaching the child to sit in the correct place and position for sufficient time to complete bladder or bowel emptying (with support or adaptations if required), by gradually increasing the time and frequency of potty/toilet sits.
- Games and appropriate communication to increase the child's understanding of what is expected.
- Appropriate use of praise and rewards for desired behaviour around toileting.
- Teaching the child to manipulate their clothing and wash their hands.

- Introduction of timed toileting when the child is most likely to need to pass urine or open their bowels, informed by a toileting chart.
- Removal of the nappies when the child is passing about half of their voids into the potty or toilet.

The children's bladder and bowel service should be consulted if there are any concerns or the child is not progressing as expected. In line with the [National Guidance for Provision of Continence Containment Products](#) to children and young people 2021, products will only be provided to children who are at least five years old and have been fully assessed and then supported for at least six months with an individualised programme to develop the skills required for toilet training. However, every child will be considered on an individual basis and decisions will be made based on the outcome of assessment.

More resources to support toilet training are available online from [Bladder & Bowel UK](#).

TOILET SKILLS ASSESSMENT TOOL

Child's Name:

NHS Number:

Date of Birth:

Assessment 1 completed by:

Job Title:

Assessment 2 completed by:

Job Title:

Date of 1st assessment:

Date of 2nd assessment:

Date of 3rd assessment:

Assessment 3 completed by:

Job Title:

BLADDER /BOWEL MATURITY

Assess 1	Assess 2	Assess 3
-------------	-------------	-------------

Bladder function – bladder emptied

1 More than once per hour,	3	Check fluid intake – adjust if necessary. Toilet training may help. If frequency persists > aged 5 yrs and toilet trained consider assessment for OAB			
2 Between 1-2 hourly	2	Indication of developing bladder maturity – toilet training may help			
3 More than 2 hourly	0	Mature bladder – consider a toilet skills development programme			

(b) Bowel function

1 Opens bowels more than three times a day	3	Exclude/treat any underlying constipation or bowel pathology			
2 Does not always have normally formed bowel movements i.e. is subjected to constipation or diarrhoea	2	Address underlying bowel problem while commencing toilet skill development programme (check Bristol Stool Form score)			
3 Has regular normally formed bowel movements	0	Mature bowel – consider a toilet skills development programme			

(c) Night-time wetting

1 Wet most nights or every night	3	If over 5 years old and dry in the day consider referral to appropriate service			
2 Has occasional or some dry nights	2	Indication of developing bladder maturity			
3 Is usually or always dry at night	0	Mature bladder – consider a toilet skills development programme			

(d) Night-time bowel movements

1 Occur more than once per week	3	Assess for underlying constipation – treat as appropriate			
2 Never or rarely occurs	0	Mature bowel – consider a toilet skills development programme			

INDEPENDENCE / AWARENESS			Assess 1	Assess 2	Assess 3
(e) Sitting on the toilet					
1 Afraid or refuses to sit	4	Consider behaviour modification programme and OT referral			
2 Sits with or without help	2	Liaise with OT if necessary re toilet adaptation/equipment			
3 Sits without help for long enough to complete voiding	0	Continue to work on other skills required for toilet training			
(f) Going to the toilet					
1 Gives no indication of need to go to the toilet	4	Consider introducing strategies to raise awareness of wet/dry/soiled and start potty/toilet sits if not already done so			
2 Gives some indication of need to go to the toilet	2	Introduce positive reinforcement for target behaviour			
3 Sometimes goes to or asks for toilet of own accord	0	Consider timed toileting and removal of nappy in the daytime when catching about half of voids in potty/toilet			
(g) Handling clothes at toilet					
1 Cannot handle clothes at all	3	If child physically able introduce programme to encourage child to learn to manage clothing			
2 Attempts or helps to dress/undress for the toilet	2	Introduce positive reinforcement for target behaviour			
3 Pulls clothes up and down without help	0	Continue to work on other skills required for toilet training			
BEHAVIOUR					
(h) Bladder control					
1 Never or rarely passes urine on toilet/potty	3	Complete baseline wetting/soiling chart to identify voiding interval and start toilet sits at times when bladder most likely to be full (minimum 60 mins apart)			
2 Passes about half of all voids on the toilet	2	Consider removal of nappy (if worn) and continue regular toilet sits at times informed by toileting chart.			
3 Can initiate a void on request	0	Remove daytime nappy when passing about half voids in potty/toilet			
(i) Bowel control					
1 Never or rarely opens bowels on toilet/potty	3	Complete baseline wetting/soiling chart to identify frequency of bowel movements and then start toilet sitting at a time when bowel more likely to be emptied e.g. after meals			
2 Opens bowels on toilet/potty sometimes	2	Start/continue to work on skills required for toilet training			
3 Opens bowels on toilet/potty every time	0	Evidence of bowel control work on other skills for toilet training			
(j) Behaviour problems, that interfere with toileting process e.g. screams when toileted					
1 Occurs frequently, i.e. once a day or more	4	Consider liaison with LD team/CAHMS re behaviour modification programme			
2 Occurs occasionally, i.e. less than once a day	2	Consider assessment to identify 'trigger' factors for behaviour e.g. sound of hand dryer			
3 Never occurs	0	Continue to work on other skills required for toilet training			
(k) Response to basic commands, e.g. "come here",					
1 Never/ Occasionally responds to commands	4	Consider introducing 'routine/social stories' to gain co-operation			
2 Usually responds	0	Continue to work on other skills required for toilet training			

PLEASE COMPLETE IN BLACK INK

Toileting Chart

Instructions Overleaf

Name:

NHS No:.....

Service contact details








Tel:

DAY 1 Date.....








DAY 2 Date.....

DAY 3 Date.....

	Drinks		Urine	Bowels		Drinks		Urine	Bowels		Drinks		Urine	Bowels
	Type	Amount				Type	Amount				Type	Amount		
6 am														
7 am														
8 am														
9 am														
10 am														
11 am														
Midday														
1 pm														
2 pm														
3 pm														
4 pm														
5 pm														
6 pm														
7 pm														
8 pm														
9 pm														
10 pm														
11 pm														
Midnight														
1 am														
2 am														
3 am														
4 am														
5 am														
TOTAL														

<p>It is important that you complete this chart as part of the assessment of your child’s bladder and bowel health and their ability to toilet train.</p> <p>INSTRUCTIONS</p> <p>Please record</p> <ol style="list-style-type: none"> 1. Type and amount of all drinks (in mls). 2. Check your child’s nappy every hour, when they are awake, and record whether wet (W) or dry (D). This can be difficult with modern “super absorbent” nappies. We suggest that you put something inside the nappy, so that you can easily tell whether your child is wet or dry. Folded kitchen roll works well; if the kitchen roll is wet, change it, but the nappy can stay on until it will not hold any more urine. 3. If your child uses the toilet or potty successfully, put (T) in the urine column. 4. Record poos in the bowel column. <p>Try and carry on for as many days as you can. Please continue for at least four days.</p>	<p>Bristol Stool Chart</p>			
		<p>Type 1. Separate hard lumps, like nuts (hard to pass)</p>		
			<p>Type 2. Sausage-shaped but lumpy</p>	<p>Type 3. Like a sausage but with cracks on the surface</p>
			<p>Type 4. Like a sausage or snake, smooth and soft</p>	<p>Type 5. Soft blobs with clear-cut edges (passed easily)</p>
		<p>Type 6. Fluffy pieces with ragged edges, a mushy stool</p>	<p>Type 7. Watery, no solid pieces, entirely liquid stool</p>	

PLEASE COMPLETE IN BLACK INK <h3 style="text-align: center;">Frequency Volume Chart</h3> Instructions Overleaf Name:				Service contact details Tel:													
DAY 1 Date.....				DAY 2 Date.....				DAY 3 Date.....									
	Drinks		Urine	Bowels			Drinks		Urine	Bowels			Drinks		Urine	Bowels	
	Type	Amount					Type	Amount					Type	Amount			
6 am																	
7 am																	
8 am																	
9 am																	
10 am																	
11 am																	
Midday																	
1 pm																	
2 pm																	
3 pm																	
4 pm																	
5 pm																	
6 pm																	
7 pm																	
8 pm																	
9 pm																	
10 pm																	
11 pm																	
Midnight																	
1 am																	
2 am																	
3 am																	
4 am																	
5 am																	
TOTAL																	

<p style="text-align: center;">INSTRUCTIONS</p> <p>Please record</p>	<p style="text-align: center;">BRISTOL STOOL CHART</p>	
<p>1. Type and amount of all drinks (in mls)</p> <p>2. The amount of urine passed in mls (measure in a jug)</p> <p>3. The time and type of bowel movements, using the Bristol Stool Chart opposite ►</p>	 <p>Type 1. Separate hard lumps, like nuts (hard to pass)</p>	 <p>Type 2. Sausage-shaped but lumpy</p>
<p>4. Any wet beds or wet clothes (write wet in the urine column). If wetting occurs estimate the amount by writing WS for a small amount WM for a medium amount WL for a large amount</p> <p>5. Indicate bedtime by writing B in the urine column</p>	 <p>Type 3. Like a sausage but with cracks on the surface</p>	 <p>Type 4. Like a sausage or snake, smooth and soft</p>
<p>6. Indicate time of waking by writing M in the urine column</p>	 <p>Type 5. Soft blobs with clear-cut edges (passed easily)</p>	 <p>Type 6. Fluffy pieces with ragged edges, a mushy stool</p>
		 <p>Type 7. Watery, no solid pieces, entirely liquid</p>

Children's Bladder and Bowel Initial Assessment Tool for all children 0 – 19 years old
(Including those with additional needs)

Child's Name	Date of birth
	NHS No:
Initial Assessment Completed by:	Presenting problem:
Contact details:	Date

Prior to undertaking the assessment the child and family should complete a bladder diary for 48 hours and bowel and night wetting diary for one week using standard documentation. Include:

- Fluid intake (what, when and how much the child has drunk).
- Frequency & consistency of bowel movements (use Bristol Stool Form chart)
Expected frequency of no more than x3 per day / no less than x3 per week
- Any soiling including time, amount, location
- Number of voids including any wetting (normal range 4 – 7 voids per day)
- Volume of voids (Expected bladder capacity = age x 30 + 30)
- Any bedwetting with estimated size of wet patch and time if known

Fluid intake (refer to chart for age-appropriate intake):

	YES	NO	ACTION
Good fluid intake: drinks 6-8 water-based drinks per day (total appropriate for age)			If no advise to adjust intake accordingly
Poor fluid intake (less than 80% of expected for age) and/or includes fizzy and caffeinated drinks			If yes advise to adjust fluid as necessary
Drinks spread evenly throughout the day?			If no advise re regular drinks including three drinks in school and last drink an hour before bed

Bowel Function:

Red Flags	YES	NO	ACTION
Any delay in passage of meconium (>48 hrs)			If yes refer infant directly to paediatrician, discuss older child with children's bladder and bowel service
Symptoms apparent within first few weeks of life			If yes refer infant directly to paediatrician. discuss older child with children's bladder and bowel service
Passing ribbon (very narrow) stools from birth			If yes refer directly to paediatrician
Concern re abdominal distension with vomiting			If yes refer directly to paediatrician
Recent leg weakness noticed			If yes refer to paediatrician

History			
Less than 3 bowel movements / week (in non breast fed baby or weaned child)			If yes consider constipation – refer to pathway
Has frequent daily soiling?			If yes consider faecal impaction – refer to constipation pathway
Stool consistency (use Bristol Stool Form Chart) reported to be 1-3 or 6-7			If yes consider potential for constipation – refer to pathway
Often or occasionally opens bowels during sleep?			If yes consider if toilet refusal in the day (possible behavioural issue) or if underlying constipation
Struggles to open bowels, withholds, has pain with bowel motions, has frequent abdominal pain?			If yes suggestive of constipation – refer to pathway
Other? (describe)			If concerned discuss with children's bladder and bowel service, or refer on to GP or paediatrician, as appropriate

Daytime Bladder Problems:

Red Flags	YES	NO	
History of repeated UTIs			If yes refer to GP for further investigation
Child (particularly girls) reported to be always wet during day			If yes refer to GP for further investigation
Any reported straining to void or weak stream			If yes refer to GP for further investigation
History			
Voids either > 7 or < 4 times per day			If yes check fluid intake to ensure within recommended amount and refer to daytime wetting pathway
Is toilet trained and has urinary incontinence during the day			If yes refer to daytime wetting pathway
Some reported frequency (voids > x7) or urgency (has to dash to the toilet)			Advise re regular toileting (e.g. 2 hourly) plus regular drinks
Child has not achieved day time dryness at all by age 3 years			If yes refer to toilet training pathway
Other? (describe)			If concerned discuss with children's bladder and bowel service

Toileting issues (from age 2 years including those with additional needs):

	YES	NO	
Behavioural problems or anxieties about using the toilet?			Consider behavioural support techniques
Has mobility or sensory problem interfering with ability to sit on toilet safely?			Consider referral to OT
Gives no indication of needing to use toilet?			If yes refer to toilet training pathway

Never or rarely passes urine or opens bowels on the toilet/potty?			If yes refer to toilet training pathway
Insists on nappy for opening bowels or other toilet refusal issue?			If yes consider behaviour modification programme
Other? (Describe)			If concerned refer to toilet training pathway and discuss with continence nurse

Night time wetting (children over the age of 4 yrs):

Red Flags	YES	NO	
Reported weight loss or excessive thirst			Refer to GP for investigation (e.g. urinalysis and blood sugar)
Concern re parental intolerance / safeguarding issues			If yes follow local safeguarding policy
History			
Is wet more than two nights a week?			If yes clinically significant refer to bedwetting pathway
Wakes after wetting			Possible overactive bladder – confirm no daytime symptoms
Other? (Describe)			Refer to bedwetting pathway and discuss with children's bladder and bowel service if concerned

N.B. ensure additional information is documented in child's notes and included on any referrals.

OUTCOME:

Advice offered: (provide details)	
Information sheets provided to family (provide details)	
Commenced on pathway: (details of pathway)	
Date for reassessment / review:	
Referred to children's bladder and bowel service	date
Signature	date

References and Resources

NICE 2010 Constipation in children and young people: diagnosis and management
<https://www.nice.org.uk/guidance/cg99>

NICE 2014 Constipation in children and young people – Quality Standard
<https://www.nice.org.uk/guidance/qs62>

NICE 2010 Bedwetting in children and young people – Guidance
<https://www.nice.org.uk/guidance/cg111>

NICE 2014 Bedwetting in children and young people – Quality Standard
<https://www.nice.org.uk/guidance/qs70>

Birth to Five HSCNI
<http://www.publichealth.hscni.net/publications/birth-five>

Healthy child programme 0 to 19: health visitor and school nurse commissioning
<https://www.gov.uk/government/publications/healthy-child-programme-0-to-19-health-visitor-and-school-nurse-commissioning>

Resources for families:

There are a range of [resources for families](#) available on the Bladder & Bowel UK website

Resources for professionals:

[Guidance for the provision of continence containment products to children and young people](#) A consensus document Bladder & Bowel UK 2021

[Managing Bladder and Bowel Issues in Nurseries, Schools and Colleges](#) Guidance for school leaders, proprietors, governors, staff and practitioners Bladder & Bowel UK and ERIC 2022

There are other [resources for professionals](#) on the Bladder & Bowel UK website

NICE has a range of useful [resources](#).